

Research Report

S E R I E S

METHAMPHETAMINE *Abuse and Addiction*

The abuse of methamphetamine — a potent psychostimulant — is an extremely serious and growing problem. Although the drug was first used primarily in selected urban areas in the Southwestern part of the United States, we now are seeing high levels of methamphetamine abuse in many areas of the Midwest, in both urban and rural settings, and by very diverse segments of the population.

As part of our Methamphetamine Research Initiative, the National Institute on Drug Abuse (NIDA) has developed this publication to provide an overview of the latest scientific findings on this drug. Methamphetamine is a powerfully addictive stimulant associated with serious health conditions, including memory loss, aggression, psychotic behavior, and potential heart and brain damage; it also contributes to increased transmission of hepatitis and HIV/AIDS.

One of NIDA's most important goals is to translate what scientists learn from research to help the public better understand drug abuse and addiction and to develop more effective strategies for their prevention and treatment. We hope this compilation of scientific information on methamphetamine will help inform readers about the harmful effects of methamphetamine abuse and will assist in prevention and treatment efforts.


Alan I. Leshner, Ph.D.
Director
National Institute on Drug Abuse

What is methamphetamine?

Methamphetamine is a powerfully addictive stimulant that dramatically affects the central nervous

system. The drug is made easily in clandestine laboratories with relatively inexpensive over-the-counter ingredients. These factors combine to make methamphetamine a drug with high potential for widespread abuse.

Methamphetamine and amphetamine use is on the rise.



Quarterly emergency room episodes due to stimulant use were tracked from 1994 to 1996. A shortage of methamphetamine was reported by epidemiologists during the last half of 1995 accounting for the significant decrease in ER episodes.

from the director



Methamphetamine is commonly known as "speed," "meth," and "chalk." In its smoked form it is often referred to as "ice," "crystal," "crank," and "glass." It is a white, odorless, bitter-tasting crystalline powder that easily dissolves in water or alcohol. The drug was developed early in this century from its parent drug, amphetamine, and was used originally in nasal decongestants and bronchial inhalers.

Methamphetamine's chemical structure is similar to that of amphetamine, but it has more pronounced effects on the central nervous system. Like amphetamine, it causes increased activity, decreased appetite, and a general sense of well-being. The effects of methamphetamine can last 6 to 8 hours. After the initial "rush," there is typically a state of high agitation that in some individuals can lead to violent behavior.

Methamphetamine is a Schedule II stimulant, which means it has a high potential for abuse and is available only through a prescription that cannot be refilled. There are a few accepted medical reasons for its use, such as the treatment of narcolepsy, attention deficit disorder, and—for short-term use—obesity; but these medical uses are limited.

What is the scope of methamphetamine abuse in the United States?

Methamphetamine abuse, long reported as the dominant drug problem in the San Diego, CA, area, has become


a substantial drug problem in other sections of the West and Southwest, as well. There are indications that it is spreading to other areas of the country, including both rural and urban sections of the South and Midwest. Methamphetamine, traditionally associated with white, male, blue-collar workers, is being used by more diverse population groups that change over time and differ by geographic area.

According to the 1996 National Household Survey on Drug Abuse, an estimated 4.9 million people (2.3 percent of the population) have tried methamphetamine at some time in their lives. In 1994, the estimate was 3.8 million (1.8 percent), and in 1995 it was 4.7 million (2.2 percent).

Data from the 1996 Drug Abuse Warning Network (DAWN), which collects information on drug-related episodes from hospital emergency departments in 21 metropolitan areas, reported that methamphetamine-related episodes decreased by 39 percent between 1994 and 1996, after a 237 percent increase between 1990 and 1994. There was a statistically significant decrease in methamphetamine-related episodes between 1995 (16,200) and 1996 (10,800). However, there was a significant increase of 71 percent between the first half of 1996 and the second half of 1996 (from 4,000 to 6,800).

NIDA's Community Epidemiology Work Group (CEWG), an early warning network of researchers that provides

The preferred method of taking methamphetamine varies among geographical regions.



Note: Calendar year in Hawaii and San Diego; State fiscal year in San Francisco
 Source: Community Epidemiology Work Group, NIDA 1997

information about the nature and patterns of drug use in major cities, reported in its June 1997 publication that methamphetamine continues to be a problem in Hawaii and in major Western cities, such as San Francisco, Denver, and Los Angeles. Increased methamphetamine availability and production are being reported in diverse areas of the country, particularly rural areas, prompting concern about more widespread use.

Drug abuse treatment admissions reported by the CEWG in December 1996 showed that methamphetamine remained the leading drug of abuse among treatment clients in the San Diego area and was second only to marijuana in Hawaii. Stimulants, including methamphetamine, accounted for smaller percentages of treatment admis-

sions in other states and metropolitan areas of the West (e.g., 5 percent in Los Angeles and Seattle and 4 percent in Texas and San Francisco). By comparison, stimulants were the primary drugs of abuse in less than 1 percent of treatment admissions in most Eastern and Midwestern metropolitan areas, except in Minneapolis-St. Paul and St. Louis, where they accounted for approximately 2 percent of total admissions.

How is methamphetamine used?


Methamphetamine comes in many forms and can be smoked, snorted, orally ingested, or injected. The drug

alters moods in different ways, depending on how it is taken.

Immediately after smoking the drug or injecting it intravenously, the user experiences an intense rush or "flash" that lasts only a few minutes and is described as extremely pleasurable. Snorting or oral ingestion produces euphoria—a high but not an intense rush. Snorting produces effects within 3 to 5 minutes, and oral ingestion produces effects within 15 to 20 minutes.

As with similar stimulants, methamphetamine most often is used in a "binge and crash" pattern. Because tolerance for methamphetamine occurs within minutes—meaning that the pleasurable effects disappear even before the drug concentration in the blood falls significantly—users try to maintain the high by

The brain – Dopamine plays an important role in the regulation of pleasure. In addition to other regions, dopamine is manufactured in nerve cells within the ventral tegmental area and is released in the nucleus accumbens and the frontal cortex.



binging on the drug.

In the 1980's, "ice," a smokeable form of methamphetamine, came into use. Ice is a large, usually clear crystal of high purity that is smoked in a glass pipe like crack cocaine. The smoke is odorless, leaves a residue that can be resmoked, and produces effects that may continue for 12 hours or more.

What are the immediate (short-term) effects of methamphetamine abuse?

As a powerful stimulant, methamphetamine, even in small doses, can increase wakefulness and physical activity and decrease appetite. A brief, intense sensation, or rush, is reported by those who smoke or inject methamphetamine. Oral ingestion or snorting produces a long-lasting high instead of a rush, which reportedly can continue for as long as half a day. Both the rush and the high are believed to result from the release of very high levels of the neurotransmitter dopamine into areas of the brain that regulate feelings of pleasure.

Methamphetamine has toxic effects. In animals, a single high dose of the drug has been shown to damage nerve terminals in the dopamine-containing regions of the brain. The large release of dopamine produced by methamphetamine is thought to con-

tribute to the drug's toxic effects on nerve terminals in the brain. High doses can elevate body temperature to dangerous, sometimes lethal, levels, as well as cause convulsions.

Short-term effects can include:

Increased attention and decreased fatigue

Increased activity

Decreased appetite

Euphoria and rush

Increased respiration

Hyperthermia

What are the long-term effects of methamphetamine abuse?

Long-term methamphetamine abuse results in many damaging effects, including addiction. Addiction is a chronic, relapsing disease, characterized by compulsive drug-seeking and drug use which is accompanied by functional and molecular changes in the brain. In addition to being addicted to methamphetamine, chronic methamphetamine abusers exhibit symptoms that can include violent behavior, anxiety, confusion, and insomnia. They also can display a number of psychotic features, including

paranoia, auditory hallucinations, mood disturbances, and delusions (for example, the sensation of insects creeping on the skin, called "formication"). The paranoia can result in homicidal as well as suicidal thoughts.

With chronic use, tolerance for methamphetamine can develop. In an effort to intensify the desired effects, users may take higher doses of the drug, take it more frequently, or change their method of drug intake. In some

Long-term effects can include:

Dependence and addiction psychosis

- paranoia*
- hallucinations*
- mood disturbances*
- repetitive motor activity*

Stroke

Weight loss

cases, abusers forego food and sleep while indulging in a form of binging known as a "run," injecting as much as a gram of the drug every 2 to 3 hours over several days until the user runs out of the drug or is too disorganized to continue. Chronic abuse can lead to psychotic behavior, characterized by intense paranoia, visual and auditory hallucinations, and out-of-control rages that can be coupled with extremely violent behavior.

Although there are no physical manifestations of a withdrawal

syndrome when methamphetamine use is stopped, there are several symptoms that occur when a chronic user stops taking the drug. These include depression, anxiety, fatigue, paranoia, aggression, and an intense craving for the drug.

In scientific studies examining the consequences of long-term methamphetamine exposure in animals, concern has arisen over its toxic effects on the brain. Researchers have reported that as much as 50 percent of the dopamine-producing cells in the brain can be damaged after prolonged exposure to relatively low levels of methamphetamine. Researchers also have found that serotonin-containing nerve cells may be damaged even more extensively. Whether this toxicity is related to the psychosis seen in some long-term methamphetamine abusers is still an open question.

How is methamphetamine different from other stimulants, like cocaine?

Methamphetamine is classified as a psychostimulant as are such other drugs of abuse as amphetamine and cocaine. We know that methamphetamine is structurally similar to amphetamine and the neurotransmitter dopamine, but it is quite different from cocaine. Although these stimulants have similar behavioral and physiological effects, there are some major differences in the basic mechanisms of how they work at the level of the nerve cell. However, the bottom line is that methamphetamine, like cocaine, results in an accumulation of the neurotransmitter dopamine, and this excessive dopamine concentration appears to produce the stimula-

tion and feelings of euphoria experienced by the user. In contrast to cocaine, which is quickly removed and almost completely metabolized in the body, methamphetamine has a much longer duration of action and a larger percentage of the drug remains unchanged in the body. This results in methamphetamine being present in the brain longer, which ultimately leads to prolonged stimulant effects.

What are the medical complications of methamphetamine abuse?

Methamphetamine can cause a variety of cardiovascular problems. These include rapid heart rate, irregular heartbeat, increased blood pressure, and irreversible, stroke-producing damage to small blood vessels in the brain.

Hyperthermia (elevated body temperature) and convulsions occur with methamphetamine overdoses, and if not treated immediately, can result in death.

Chronic methamphetamine abuse can result in inflammation of the heart lining, and among users who inject the drug, damaged blood vessels and skin abscesses. Methamphetamine abusers also can have episodes of violent behavior, paranoia, anxiety, confusion, and insomnia. Heavy users also show progressive social and occupational deterioration. Psychotic symptoms

Although both methamphetamine and cocaine are psychostimulants, there are differences between them.

Methamphetamine	vs.	Cocaine
<i>Man-made</i>		<i>Plant-derived</i>
<i>Smoking produces a high that lasts 8-24 hours</i>		<i>Smoking produces a high that lasts 20-30 minutes</i>
<i>50% of the drug is removed from the body in 12 hours</i>		<i>50% of the drug is removed from the body in 1 hour</i>
<i>Limited medical use</i>		<i>Used as a local anesthetic in some surgical procedures</i>

can sometimes persist for months or years after use has ceased.

Acute lead poisoning is another potential risk for methamphetamine abusers. A common method of illegal methamphetamine production uses lead acetate as a reagent. Production errors may therefore result in methamphetamine contaminated with lead. There have been documented cases of acute lead poisoning in intravenous methamphetamine abusers.

Fetal exposure to methamphetamine also is a significant problem in the United States. At present, research indicates that methamphetamine abuse during pregnancy may result in prenatal complications, increased rates of premature delivery, and altered neonatal behavioral patterns, such as abnormal reflexes and extreme irritability. Methamphetamine abuse during pregnancy may be linked also to congenital deformities.

Are methamphetamine abusers at risk for contracting HIV/AIDS and hepatitis B and C?

Increased HIV and hepatitis B and C transmission are likely consequences of increased methamphetamine abuse, particularly in individuals who inject the drug and share injection equipment. Infection with HIV and other infectious diseases is spread

among injection drug users primarily through the reuse of contaminated syringes, needles, or other paraphernalia by more than one person. In nearly one-third of Americans infected with HIV, injection drug use is a risk factor, making drug abuse the fastest growing vector for the spread of HIV in the nation.

Research also indicates that methamphetamine and related psychomotor stimulants can increase the libido in users, in contrast to opiates which actually decrease the libido. However, long-term methamphetamine use may be associated with decreased sexual functioning, at least in men. Additionally, methamphetamine seems to be associated with rougher sex which may lead to bleeding and abrasions. The combination of injection and sexual risks may result in HIV becoming a greater problem among methamphetamine abusers than among opiate and other drug abusers, something that already seems to be occurring in California.

NIDA-funded research has found that, through drug abuse treatment, prevention, and community-based outreach programs, drug abusers can change their HIV risk behaviors. Drug use can be eliminated and drug-related risk behaviors, such as needle-sharing and unsafe sexual practices, can be reduced significantly thus decreasing the risk of exposure. Therefore, drug abuse treatment is also highly effective in preventing the spread of HIV, hepatitis B, and hepatitis C.

What treatments are effective for methamphetamine abusers?

At this time the most effective treatments for methamphetamine addiction are cognitive behavioral interventions. These approaches are designed to help modify the patient's thinking, expectancies, and behaviors and to increase skills in coping with various life stressors. Methamphetamine recovery support groups also appear to be effective adjuncts to behavioral interventions that can lead to long-term drug-free recovery.

There are currently no particular pharmacological treatments for dependence on amphetamine or amphetamine-like drugs such as methamphetamine. The current pharmacological approach is borrowed from experience with treatment of cocaine dependence. Unfortunately, this approach has not met with much success since no single agent has proven efficacious in controlled clinical studies. Antidepressant medications are helpful in combating the depressive symptoms frequently seen in methamphetamine users who recently have become abstinent.

There are some established protocols that emergency room physicians use to treat individuals who have had a methamphetamine overdose. Because hyperthermia and convulsions are common and often fatal complications of such overdoses, emer-

gency room treatment focuses on the immediate physical symptoms. Overdose patients are cooled off in ice baths, and anti-convulsant drugs may be administered also.

Acute methamphetamine intoxication can often be handled by observation in a safe, quiet environment. In cases of extreme excitement or panic, treatment with anti-anxiety agents such as benzodiazepines has been helpful, and in cases of methamphetamine-induced psychoses, short-term use of neuropeptides has proven successful.

Where can I get further scientific information about methamphetamine abuse?

To learn more about methamphetamine and other drugs of abuse, contact the National Clearinghouse for Alcohol and Drug Information (NCADI) at 1-800-729-6686. Information specialists are available to assist you in locating needed information and resources. Information can be accessed also through the NIDA World Wide Web site (<http://www.nida.nih.gov>) or the NCADI Web site (<http://www.health.org>).

Fact sheets on health effects of drug abuse and other topics can be ordered free of charge, in English and Spanish, by calling NIDA INFOFAX at 1-800-NIH-NIDA (1-800-644-6432) or

1-888-TTY-NIDA (1-888-889-6432) for the hearing impaired.

A list of fact sheet topics follows:

Health Effects of Specific Drugs

- Cigarettes and other tobacco products
- Crack and cocaine
- Ecstasy
- Heroin
- Inhalants
- LSD
- Marijuana
- Methamphetamine
- Pain medications
- PCP
- Ritalin
- Rohypnol and GHB
- Steroids (anabolic)

Drug Abuse and AIDS Lessons from Prevention Research

Treatment Research

- Treatment methods
- Treatment medications
- Treatment methods for women
- Behavior change through treatment

Trends and Surveys

- Costs to society from drug abuse
- High school and youth trends
- Hospital visits and deaths
- Nationwide trends
- Pregnancy and drug abuse trends
- Treatment trends
- Workplace trends

News Releases on Research Findings

Information about NIDA

- Home page on the World Wide Web
- Mission and structure

- Opportunities for special populations
- Funding opportunities
- Upcoming events and conferences

Other Drug Abuse Information Services

NIDA web site
<http://www.nida.nih.gov>

Access information on the internet

- What's new on the NIDA web site
- Information on drugs of abuse
- Publications and communications (including *NIDA Notes*)
- Calendar of events
- Links to NIDA organizational units
- Funding information (including program announcements and deadlines)
- International activities
- Links to related web sites (access to web sites of many other organizations in the field)

NCADI web site
<http://www.health.org>

Glossary

Addiction: a chronic, relapsing disease, characterized by compulsive drug-seeking and drug use and by neurochemical and molecular changes in the brain.

Analog: a chemical compound that is similar to another drug in its effects but differs slightly in its chemical structure.

Benzodiazepines: drugs that relieve anxiety or are prescribed as sedatives; among the most widely prescribed medications, including valium and librium.

Central nervous system (CNS): the brain and spinal cord.

Craving: a powerful, often uncontrollable desire for drugs.

Designer drug: an analog of a restricted drug that has psychoactive properties.

Detoxification: a process of allowing the body to rid itself of a drug while managing the symptoms of withdrawal; often the first step in a drug treatment program.

Dopamine: a neurotransmitter present in regions of the brain that regulate movement, emotion, motivation, and feelings of pleasure.

Narcolepsy: a disorder characterized by uncontrollable attacks of deep sleep.

Physical dependence: an adaptive physiological state that occurs with regular drug use and results in a withdrawal syndrome when drug use stops.

Psychosis: a mental disorder characterized by symptoms such as delusions or hallucinations that indicate an impaired conception of reality.

Rush: a surge of euphoric pleasure that rapidly follows administration of a drug.

Serotonin: a neurotransmitter that has been implicated in states of consciousness, mood, depression, and anxiety.

Tolerance: a condition in which higher doses of a drug are required to produce the same effect as experienced initially; often leads to physical dependence.

Toxic: temporary or permanent drug effects that are detrimental to the functioning of an organ or group of organs.

Withdrawal: a variety of symptoms that occur after use of an addictive drug is reduced or stopped.

Resources

"Blood Level of Intravenous Drug Users," by R.L. Norton, B.T. Burton, and J. McGirr. *Journal of Clinical Toxicology* 34(4):425-30, 1996.

Epidemiologic Trends in Drug Abuse: Vol. 1. Highlights and Executive Summary of the Community Epidemiology Work Group, June 1997. NIH Pub. No. 98-4207. National Institute on Drug Abuse, 1997.

Epidemiologic Trends in Drug Abuse: Vol. 1. Highlights and Executive Summary of the Community Epidemiology Work Group, December 1996. NIH Pub. No. 97-4204. National Institute on Drug Abuse, 1997.

"Integrating Treatments for Methamphetamine Abuse: A Psychosocial Perspective," by A. Huber, W. Ling, S. Shoptaw, V. Gulati, P. Brethen, and R. Rawson. *Journal of Addictive Diseases*, 16(4):41-50, 1997.

"Like Methamphetamine, Ecstasy May Cause Long-Term Brain Damage," by R. Mathias. *NIDA Notes* 11:7, 1996.

Methamphetamine Abuse (NIDA Capsules). National Institute on Drug Abuse, September 1997.

National Methamphetamine Strategy. U.S. Department of Justice, 1996.

National Survey Results on Drug Use From the Monitoring the Future Study, 1975-1994, Vol. I: Secondary School Students. NIH Pub. No. 93-3498. National Institute on Drug Abuse, 1995.

National Survey Results on Drug Use From the Monitoring the Future Study, 1975-1994, Vol. II: College Students and Young Adults. NIH Pub. No. 96-4027. National Institute on Drug Abuse, 1995.

"NIDA Survey Provides First National Data on Drug Abuse During Pregnancy," by R. Mathias. *NIDA Notes* 10:6-7, 1995.

"Preliminary Results from the 1996 National Household Survey on Drug Abuse". Substance Abuse and Mental Health Services Administration, 1997.

Year-End Preliminary Estimates from the 1996 Drug Abuse Warning Network. Substance Abuse and Mental Health Services Administration, November 1997.

NIDA
NATIONAL INSTITUTE
ON DRUG ABUSE

NIH Publication Number 98-4210.
Printed April 1998.
Feel free to reprint this publication.