

## Improvements to the Synthesis of Psilocybin and a Facile Method for Preparing the *O*-Acetyl Prodrug of Psilocin

David E. Nichols,\* Stewart Frescas

Department of Medicinal Chemistry and Molecular Pharmacology, School of Pharmacy and Pharmacal Sciences, Purdue University, West Lafayette, Indiana 47907, USA

Fax +1(765)4941414; E-mail drdave@pharmacy.purdue.edu

Received 3 December 1998; revised 11 February 1999

**Abstract:** An improved procedure to accomplish the *O*-phosphorylation of 4-hydroxy-*N,N*-dimethyltryptamine (psilocin **5**) is reported that utilizes reaction between the *O*-lithium salt of **5** and tetra-*O*-benzylpyrophosphate. The *O*-benzyl groups were removed by catalytic hydrogenation over palladium on carbon to afford *N,N*-dimethyl-4-phosphoryloxytryptamine (psilocybin, **1**). In view of difficulties encountered in the preparation of **1**, it is suggested that 4-acetoxy-*N,N*-dimethyltryptamine (**2**) may be a useful alternative for pharmacological studies. The latter was obtained following catalytic *O*-debenzylation of 4-benzyloxy-*N,N*-dimethyltryptamine in the presence of acetic anhydride and sodium acetate.

**Key words:** psilocin, psilocybin, tetra-*O*-benzylpyrophosphate, phosphorylation

Recently, several laboratories have initiated clinical studies of hallucinogenic (psychedelic) agents.<sup>1–3</sup> This renewed interest suggests that there may be some demand for investigational substances that are suitably pure for human use that can be prepared in a relatively economical fashion. Hallucinogens are not commercially available in large quantities or in purities suitable for human studies, and research will likely be carried out only with drugs produced by custom synthesis. Of the various drugs that might be of interest for this work, most of them, including mescaline, LSD, DMT, and various substituted amphetamines are synthesized relatively easily. Indeed, many hallucinogens are routinely manufactured in clandestine laboratories.

By contrast, the synthesis of psilocybin, *N,N*-dimethyl-4-phosphoryloxytryptamine (**1**), is more challenging. Nevertheless, psilocybin has pharmacological features that make it attractive for clinical research, including a relatively short duration of action. The increasing worldwide popularity of psilocybin-containing mushrooms as recreational drugs also points to the need for more research with psilocybin.


We re-examined the synthesis of psilocybin reported by Hofmann and co-workers.<sup>4</sup> Although their approach still remains useful, there were several weak points that could be addressed to improve the yields and purities of the final compound.

The overall synthetic route is shown in Scheme 1. The most troublesome step is the last, the phosphorylation of psilocin. In the original synthesis by Hofmann et al.,<sup>4</sup> the phosphorylation step was accomplished using *O,O*-dibenzylphosphoryl chloride, an unstable reagent that was used without purification as a solution in carbon tetrachloride. Furthermore, the final yield of psilocybin was less than 20%. In view of the overall difficulty in preparing this material and its precursors, such a low yield in the last step was deemed unacceptable.


The present synthesis employs a phosphorylation step using tetrabenzylpyrophosphate, a stable, crystalline reagent. The phosphorylation step was complicated by the previously unreported *extremely* labile nature of the *O,O*-dibenzyl ester of psilocybin. Hydrolytic cleavage of one of the *O*-benzyl groups occurred rapidly in the presence of water, at room temperature, and neutral pH. The purification of the resulting zwitterionic material was much more complicated than for the basic *O,O*-dibenzyl material.

Illustrated in Scheme 2 is the facile preparation of 4-acetoxy-DMT<sup>5</sup> **2**. This *O*-acetyl prodrug of psilocin is much more easily prepared than psilocybin, and may offer an economical alternative for clinicians wishing to study the psychopharmacology of psilocin. This material is readily crystallized as the fumarate salt, and is considerably more stable than psilocin itself. It would seem to be an ideal prodrug to replace psilocybin in future clinical studies, since psilocin is the principal metabolite of psilocybin.<sup>6</sup>


The classical Speeter and Anthony synthesis of tryptamines from indoles served as the precedent for this work.<sup>7</sup> The key reaction of oxalyl chloride with 4-benzyloxyindole was, however, sluggish. Similarly, the reduction of the 4-substituted glyoxalylamide **3** was much slower than for indoles without substitution at this position. TLC was used to monitor the complete disappearance of starting material and intermediate reduction products. The *O*-benzyl group was then readily removed by catalytic hydrogenolysis to afford 4-hydroxy-*N,N*-dimethyltryptamine (psilocin; **5**).



Psilocybin (**1**)



4-Acetoxy-DMT (**2**)




Scheme 1

After experimentation with a variety of phosphorylating agents, it was finally decided that tetrabenzylpyrophosphate (TBPP) was the most suitable reagent.<sup>8</sup> This crystalline and stable material is commercially available (Aldrich), but can also be synthesized readily on a multi-gram scale.

The most convenient base for the phosphorylation step proved to be butyllithium. Generation of the lithium salt of psilocin in THF at  $-70^\circ\text{C}$ , followed by addition of 1.1 equivalents of TBPP, led to the *O,O*-dibenzyl ester of psilocybin, with the generation of one equivalent of lithium *O,O*-dibenzylphosphate that must ultimately be removed. While ordinarily removal of the lithium salt would not be problematic, washing the organic reaction mixture with water led to unexpected and rapid hydrolysis of one of the *O*-benzyl groups. Judicious exclusion of traces of water allowed the isolation of *O,O*-dibenzyl ester that was nearly free of 6. The *O,O*-dibenzyl intermediate proved to be so sensitive to water, however, that it was more practical to use an aqueous workup, allow hydro-

lysis to occur, and isolate a product that was largely the zwitterionic *O*-monobenzylphosphate 6.

Catalytic hydrogenolysis of the crude *O*-benzyl ester led to the formation of psilocybin (1). The procedure was complicated by small amounts of phosphoric acid generated from residual dibenzylphosphoric acid carried from the previous step into the hydrogenolysis reaction. This highly acidic material leads to discoloration of the product and prevents satisfactory crystallization. The problem was solved through the use of anion exchange resin to titrate the phosphoric acid. The reported pH of a solution of psilocybin in 50% aqueous ethanol is 5.2.<sup>9</sup> Anion exchange resin ( $-\text{OH}$  form) was added in portions, with vigorous and extended stirring, to the filtered reaction solution until the pH of the solution was 5.2. When this pH was reached, the resin was removed by filtration and the filtrate was concentrated under vacuum. The crude product was then recrystallized from a small amount of methanol, and a large volume of isopropanol, followed by



Scheme 2

storage in the freezer. Psilocybin (**1**) crystallized as long colorless needles.

As a potential replacement for **1**, 4-Acetoxy-*N,N*-dimethyltryptamine (**2**) fumarate was conveniently prepared by shaking under hydrogen a mixture of **4**, acetic anhydride, and sodium acetate in benzene with Pd/C in a Parr low pressure hydrogenation apparatus. Following uptake of the required amount of hydrogen corresponding to *O*-debenzylation, the catalyst and insoluble salts were removed by filtration. One molar equivalent of fumaric acid was added to the filtrate, and the solution was concentrated to dryness under vacuum. The resulting solid was recrystallized to afford white crystals of the desired product. This material was stable when stored in the cold, but slowly darkened on storage for several months at ambient temperature.

Mps were determined on a Thomas-Hoover Meltemp melting point apparatus and are uncorrected except where indicated. <sup>1</sup>H NMR spectra were recorded on a Bruker ARX 300 MHz spectrometer. Chemical shifts are reported in  $\delta$  values (ppm) relative to an internal standard of TMS in CDCl<sub>3</sub>, except where noted. Abbreviations used in NMR analysis are as follows: s, singlet; d, doublet; t, triplet; m, multiplet; br s, broad singlet; dd, doublet of doublets, dt, doublet of triplets. Microanalyses were obtained from the Purdue Microanalytical Laboratory. A low pressure Parr apparatus was used for all hydrogenations. Solvents and reagents were used as purchased, except as noted. THF was distilled from potassium metal/benzophenone ketyl. All other compounds were purchased from commercial sources.

#### 4-Benzoyloxyindol-3-yl-*N,N*-dimethylglyoxylamide (**3**)

A solution of 4-benzoyloxyindole (17.5 g, 0.078 mol) (Biosynth) in anhyd Et<sub>2</sub>O (500 mL) was mechanically stirred in a 1 L, 3 necked flask and cooled in an ice-salt bath to an internal temperature of 0 °C. Oxalyl chloride (20.3 g, 0.16 moles) was added dropwise at a rate that maintained an internal temperature between 0–5 °C. Stirring was continued for 3 h at a temperature between 5–10 °C with a gentle argon sparge to remove evolved HCl. The argon sparge was replaced by a gas inlet tube and a dry ice/acetone condenser. Anhyd dimethylamine was then bubbled into the reaction with cooling and vigorous stirring until a pH (determined by moist pH paper) between 9 and 11 was achieved. At this time, the orange color of the initial solution had been mostly discharged, and the reaction had the appearance of a slightly off-white slurry with a few flecks of yellow unreacted starting material. CH<sub>2</sub>Cl<sub>2</sub> (20 mL) was added to assist solubilization of the unreacted material and the reaction was stirred for an additional 6 h to yield finally an off-white slurry. Et<sub>2</sub>O (150 mL) was added, and the mixture was cooled to 10 °C. The white solids were collected by suction filtration on filter paper in a Buchner funnel and then were suspended in distilled H<sub>2</sub>O (250 mL) and stirred for 1 h to remove dimethylamine hydrochloride. The slurry was filtered, and the collected solids were washed on the filter with distilled H<sub>2</sub>O (3 x 75 mL) and hexane (75 mL) and dried overnight in a vacuum oven. The dried product weighed 18.3 g. The organic filtrates and washes were combined and the solvent was removed by rotary evaporation. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (100 mL) and the organic solution was washed with distilled H<sub>2</sub>O (2 x 50 mL) and brine (2 x 50 mL). After drying (MgSO<sub>4</sub>) the volume was reduced by rotary evaporation. The concentrated residual solution was subjected to flash chromatography over silica gel, first eluting with CH<sub>2</sub>Cl<sub>2</sub> to recover unreacted indole (1.3 g, 7.4%), followed by elution with 10% MeOH in CH<sub>2</sub>Cl<sub>2</sub> to recover 3.3 g of **3**. The latter was combined with the initial product to provide a total weight of

21.6 g (85.9%). The crude product was recrystallized from MeOH/EtOAc to give 19.5 g (77%) of **3** with mp 152–155 °C (Lit.<sup>4</sup> mp 146–150 °C).

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>):  $\delta$  = 2.88, 2.92 (2s, 6H, NCH<sub>3</sub>), 5.21 (s, 2H, CH<sub>2</sub>), 6.60 (d, 1H, *J* = 7.92 Hz, Ar), 6.86 (d, 1H, *J* = 8.04 Hz, Ar), 7.27–7.37 (m, 3H, Ar), 7.50 (m, 3H, Ar), 10.07 (br s, 1H, NH).

#### 4-Benzoyloxy-*N,N*-dimethyltryptamine (**4**)

A slurry of LiAlH<sub>4</sub> (8.90 g, 0.234 mol) in anhyd THF (100 mL) was prepared in a 2 L, 3-neck flask, previously dried with a heat gun under an argon purge. The flask was fitted with a reflux condenser, mechanical stirrer, and addition funnel. Anhyd dioxane (200 mL) was added, and the mixture was heated to 60 °C on an oil bath. 4-benzoyloxyindol-3-yl-*N,N*-dimethylglyoxylamide (**3**) (14.5 g, 0.045 moles) was dissolved in a mixture of dioxane (250 mL) and THF (150 mL) and, with rapid stirring, this solution was added dropwise over 1 h. The oil bath temperature was held at 70 °C for 4 h, followed by vigorous reflux overnight (16 h) at an oil bath temperature of 95 °C. Thin layer chromatographic analysis (9:1 CH<sub>2</sub>Cl<sub>2</sub>/MeOH; silica plates) showed nearly complete reduction. The reaction was heated at reflux for an additional 4 h and then cooled to 20 °C. A solution of distilled H<sub>2</sub>O (27 mL) in THF (100 mL) was added dropwise, resulting in a gray flocculent precipitate. Et<sub>2</sub>O (250 mL) was added to assist breakup of the complex and improve filtration. This slurry was stirred for 1 h and the mixture was then filtered with a Buchner funnel. The filter cake was washed on the filter with warm Et<sub>2</sub>O (2 x 250 mL) and was broken up, transferred back into the reaction flask and vigorously stirred with additional hot Et<sub>2</sub>O (500 mL). This slurry was filtered, and the cake was washed on the filter with Et<sub>2</sub>O (150 mL) and hexane (2 x 150 mL). All of the organic filtrates were combined and dried (MgSO<sub>4</sub>). After the drying agent was removed by filtration, the filtrate was concentrated under vacuum at 40 °C and dried under high vacuum at 0.01 mm Hg, leading to crystallization of the residue as a white waxy solid. Recrystallization from EtOAc yielded 12.57 g, (94.8%) of **4** with mp 124–126 °C (lit.<sup>4</sup> mp 125–126 °C).

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>):  $\delta$  = 2.14 (s, 6H, NCH<sub>3</sub>), 2.58 (t, 2H, *J* = 8.0 Hz, CH<sub>2</sub>), 3.04 (t, 2H, *J* = 8.0 Hz, CH<sub>2</sub>), 5.17 (s, 2H, CH<sub>2</sub>), 6.52 (d, 1H, *J* = 7.6 Hz, Ar), 6.87 (s, 1H, Ar), 6.93 (d, 1H, *J* = 8.0 Hz, Ar), 7.04 (t, 1H, *J* = 7.9 Hz, Ar), 7.29–7.39 (m, 3H, Ar), 7.49 (br d, 2H, *J* = 7.0 Hz, Ar), 8.06 (br s, 1H, NH).

#### 4-Hydroxy-*N,N*-dimethyltryptamine (Psilocin; **5**):

A solution of **4** (4.0 g, 0.0135 moles) in 95% EtOH (250 mL) was added to 1.5 g Pd/C (10% w/w) in a 500 mL Parr low pressure hydrogenation bottle. The mixture was shaken under 60 psig of H<sub>2</sub> pressure for 2 h. The catalyst was removed by vacuum filtration through Celite and was washed on the filter with EtOH (3 x 50 mL). The filtrate was concentrated by rotary evaporation. The clear residual oil was placed under high vacuum and induced to crystallize by seeding. The white crystalline powder (2.68 g, 97.0%) was used in the next step without further purification.

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>):  $\delta$  = 2.36 (s, 6H, NCH<sub>3</sub>), 2.68 (m, 2H, CH<sub>2</sub>),<sup>10</sup> 2.93 (m, 2H, CH<sub>2</sub>),<sup>10</sup> 6.54 (d, 1H, *J* = 7.6, Ar), 6.83 (br d, 2H, *J* = 12.2 Hz, Ar), 7.03 (t, 1H, *J* = 7.8 Hz, Ar), 7.86 (br s, 1H, NH), 13.2 (br s, 1H, OH; observed only by integration).

#### 4-*O*-Monobenzylphosphoryloxy-*N,N*-dimethyltryptamine (**6**)

A solution of 0.45 g (2.2 mmol) of psilocin (**5**) and 0.073 g (0.73 mmol) of diisopropylamine in anhyd THF (50 mL) was magnetically stirred in a 100 mL 3-necked flask and was cooled to –78 °C in a dry ice-acetone bath. A 2.5 M solution (1.14 mL, 2.85 mmol) of BuLi in hexane was added dropwise using a syringe. After complete addition, the reaction was stirred for 3 min and tetrabenzylpyrophosphate<sup>8</sup> (1.50 g, 2.8 mmol) was added all at once. The dry ice-acetone bath was replaced by an ice-salt bath,

